

- 10 **Purnell JQ**, Kahn SE, Schwartz RS, *et al*. Relationship of insulin sensitivity and apo-B levels to intra-abdominal fat in subjects with familial combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 2001;**21**:567–72.
- 11 **Keulen ETP**, Voors-Pette C, de Bruin TWA. Familial dyslipidemic hypertension syndrome: familial combined hyperlipidemia, and the role of abdominal fat mass. *Am J Hypertens* 2001;**14**:357–63.
- 12 **Voors-Pette C**, de Bruin TW. Excess coronary artery disease, cardiac risk factors and central obesity in familial combined hyperlipidemia (FCHL). *Atherosclerosis* 2001;**157**:481–9.
- 13 **Biegelsen ES**, Loscalzo J. Endothelial function and atherosclerosis. *Coron Artery Dis* 1999;**10**:241–56.
- 14 **Celermajer DS**, Sorensen KE, Gooch VM, *et al*. Non-invasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet* 1992;**340**:1111–15.
- 15 **Neunteufl T**, Katzenschlager R, Hassan A, *et al*. Systemic endothelial dysfunction is related to the extent and severity of coronary artery disease. *Atherosclerosis* 1997;**129**:111–18.
- 16 **Yataco AR**, Corretti MC, Gardner AW, *et al*. Endothelial reactivity and cardiac risk factors in older patients with peripheral arterial disease. *Am J Cardiol* 1999;**83**:754–8.
- 17 **Celermajer DS**, Sorensen KE, Bull C, *et al*. Endothelium-dependent dilation in the systemic arteries of asymptomatic subjects relates to coronary risk factors and their interaction. *J Am Coll Cardiol* 1994;**24**:1468–74.
- 18 **Liguori C**, Paolillo A, Pietrosanto A. An automatic measurement system for the evaluation of carotid intima-media thickness. *IEEE Trans Instr Meas* 2001;**50**:1684–91.
- 19 **Strong JP**, Malcom GT, McMahan CA, *et al*. Prevalence and extent of atherosclerosis in adolescents and young adults. Implications for prevention from the Pathobiological Determinants of Atherosclerosis in Youth Study. *JAMA* 1999;**281**:727–35.
- 20 **Greenland P**, Abrams J, Aurigemma GP, *et al*. Prevention Conference V: beyond secondary prevention: identifying the high-risk patient for primary prevention: noninvasive tests of atherosclerotic burden: Writing Group III. *Circulation* 2000;**101**:e16–22.
- 21 **Anderson TJ**, Uehata A, Gerhard MD, *et al*. Close relation of endothelial function in the human coronary and peripheral circulations. *J Am Coll Cardiol* 1995;**26**:1235–41.
- 22 **Keulen ET**, Kruijschoop M, Schaper NC, *et al*. Increased intima-media thickness in familial combined hyperlipidemia associated with apolipoprotein B. *Arterioscler Thromb Vasc Biol* 2002;**22**:283–8.
- 23 **Ylitalo K**, Syvanne M, Salonen R, *et al*. Carotid artery intima-media thickness in Finish families with familial combined hyperlipidemia. *Atherosclerosis* 2002;**162**:171–8.
- 24 **Demacker PN**, Veerkamp MJ, Bredie SJ, *et al*. Comparison of the measurements of lipids and lipoproteins versus assay of apolipoprotein B for estimation of coronary artery disease risk: a study in familial combined hyperlipidemia. *Atherosclerosis* 2000;**153**:483–90.
- 25 **Veerkamp MJ**, de Graaf J, Bredie SJ, *et al*. Diagnosis of familial combined hyperlipidemia based on lipid phenotype expression in 32 families. Results of a 5-year follow-up study. *Arterioscler Thromb Vasc Biol* 2002;**22**:274–82.
- 26 **Veerkamp MJ**, de Graaf J, Hendriks JC, *et al*. Nomogram to diagnose familial combined hyperlipidemia on the basis of results of a 5-year follow-up study. *Circulation* 2004;**109**:2980–5.
- 27 **Sattar N**, Petrie JR, Jaap AJ. The atherogenic lipoprotein phenotype and vascular endothelial dysfunction. *Atherosclerosis* 1998;**138**:229–35.
- 28 **Steinberg HO**, Chaker H, Leaming R, *et al*. Obesity/insulin resistance is associated with endothelial dysfunction: implications for the syndrome of insulin resistance. *J Clin Invest* 1996;**97**:2601–10.
- 29 **Iannuzzi A**, Licenziati MR, Acampora C, *et al*. Preclinical changes in the mechanical properties of abdominal aorta in obese children. *Metabolism* 2004;**53**:1243–6.
- 30 **Visser M**, Bouter LM, McQuillan GM, *et al*. Elevated C-reactive protein levels in overweight and obese adults. *JAMA* 1999;**282**:2131–5.
- 31 **Rexrode KM**, Pradhan A, Manson JE, *et al*. Relationship of total and abdominal adiposity with CRP and IL-6 in women. *Ann Epidemiol* 2003;**13**:674–82.


IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2006.094656

Right ventricular pacing-induced hypotension

A cardiologist was consulted to evaluate a patient with intermittent hypotension. A 71-year-old woman with a history of coronary artery disease and coronary bypass surgery, type 2 diabetes and sick sinus syndrome s/p dual-chamber pacemaker implantation had been admitted to the intensive care unit with pulmonary oedema.

The problem was a ventricular pacing-related decrease in systolic blood pressure from 110 to 70 mm Hg. Even few ventricle-paced beats led to a significant decrease in systolic blood pressure that corrected after the return of native conduction. Programmed atrioventricular interval was short and thus led to intermittent atrial synchronous ventricular pacing. As the atrioventricular synchrony was preserved, the decrease in cardiac output was not due to the classic pacemaker syndrome. The decrease in cardiac output was most likely secondary to right



ventricular apical pacing that results in an asynchronous ventricular contraction. To avoid unnecessary right ventricular pacing, a longer atrioventricular interval was programmed and hypotensive episodes resolved (panels A and B).

J Lehtonen, S Pakarinen
jukka.lehtonen@hus.fi